




representative
Image



BEST AVAILABLE C

[Show "fr" image]

BARAN, Robert, J.;

none

(No patents reference this one)

Intellectual Property Network for Business

IPN Home | Search | Order | Shopping Cart | Login | Help



WO9709049A2: METHOD FOR EFFECTING VASODILATION WITH (1,5-INTER)ARYL PROSTAGLANDIN DERIVATIVES

[View Images \(31 pages\)](#) | [View Cart](#)
[Premium Data 1: PDF \(~2950 KB\)](#) | [TIFF \(~2330 KB\)](#) | [Fax](#) | [More choices...](#)
Inventor(s)

CHEN, June, 28771 Via Los Arboles, San Juan Capistrano, CA 92675, United States of America
 BURK, Robert, M., 1337 Cerritos Drive, Laguna Beach, CA 92651, United States of America
 WOODWARD, David, F., 23152 Tulip, Lake Forest, CA 92630, United States of America

Applicant(s)

ALLERGAN, 8301 Mars Drive, Waco, TX 76712, United States of America

Issued/Filed Dates

March 13, 1997 / Aug. 28, 1996

Application Number

WO1996US0013889

IPC Class

A61K 031/557; C07C 405/00

Priority Number(s)

Sept. 1, 1995 US4995008522775

Examined Countries

AU, CA, JP, European patent: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE

Abstract

The present invention provides a method of effecting vasodilation, comprising: administering to a warm blooded animal in need of such treatment, an effective amount of (1,5-inter)aryl prostaglandin derivative represented by formula (I), wherein n is 0 or an integer of from 1 to 6; R is selected from the group of radicals represented by the formulae CO₂R', CONR'₂, CH₂OR' and SO₂NR'₂, wherein R' is hydrogen or a lower alkyl radical having from one to six carbon atoms; R'' is hydrogen or an acyl radical having the formula: (CO)R''' wherein R''' is a saturated or unsaturated acyclic hydrocarbon radical having from 1 to about 10 carbon atoms, or -(CH₂)_mR''' wherein m is 0 or an integer of from 1 to 6 and R''' is an aliphatic ring having from 3 to 7 carbon atoms or an aryl group, e.g. phenyl, or a heteroaryl group, e.g. thienyl, furanyl or pyridyl, and preferably R''' is a lower alkyl group having from 1 to 6 carbon atoms; the hatched triangular segments represent alpha oriented bonds, the solid triangular segments represent beta oriented bonds and the wavy segments represent bonds that may be in either the cis or trans orientation. More preferably said (1,5 inter)aryl prostaglandin derivative is a compound represented by formula (II). Most preferably, said (1,5-inter)aryl prostaglandin derivative is a compound of formula (II) wherein R is CO₂R', R' is hydrogen and n is 0.

[\[Show "fr" Abstract\]](#)

BEST AVAILABLE COPY